

AGROMETEOROLOGICAL BULLETIN KENYA METEOROLOGICAL DEPARTMENT

17th Dekad, 11th to 20th June, 2009

Issue No. 17/2009, Season: MJJ

HIGHLIGHTS

- During the 17th Dekad i.e. 11th 20th June 2009, most parts of the country experienced significant reduction in total Dekadal rainfall amounts, except for the Coastal Regions which experienced heavy rainfall. Western, Nyanza Provinces and Central Rift Valley, experienced light to moderate rainfall with Kakamega, Kisii, Kisumu, Kericho, Kitale, Eldoret and Nakuru and recording 63.5, 60.9, 21.2, 50.6, 9.7, 4.9 and 1.8 mm respectively. Central Province, parts of Eastern Province bordering Mt. Kenya, Nairobi Area and its environs experienced light rainfall with Nyeri, Mwea, Embu, Dagoretti, JKIA, Wilson Airport and Kabete recording a Dekadal rainfall total of 8.1, 1.7, 8.9, 23.0, 2.6, 11.7 and 21.8 mm respectively. Coast Province experienced heavy rainfall with Mombasa, Mtwapa, Mshabaha, Malindi and Lamu recording a Dekadal rainfall totals of 43.5, 55.8, 138.4, 143.0, and 133.7mm respectively. The rest of the Country experienced mainly sunny conditions within significant light rains over few places. (Fig 1)
- Day time temperatures were relatively lower due to increased cloud cover over most parts of the country, with Lodwar, Mandera, Wajir, Garissa, Kakamega, Kisumu, Mwea, Makindu, Dagoretti and Mombasa recording a Dekadal Mean day Maximum of 35.5, 35.5, 34.2, 33.1, 27.7, 30.0, 28.1, 29.8, 22.7 and 29.1 deg. Celsius respectively. (Fig 2)
- Night temperatures were relatively lower over Western and Central Rift Valley, Central Highlands, Nairobi area and its environs due to reduced cloud cover during the Dekad with Nyahururu, Narok, Eldoret Airport, Nyeri, Kabete and Dagoretti recording a Dekadal mean minimum of 5.5, 9.2, 8.0, 13.3, 13.4 and 13.6 deg Celsius respectively. (Figs 1, 2)

- The pastoral regions and game parks of Southern Rift Valley, North Eastern, South Eastern Lowlands and the Coastal regions, experienced mainly sunny conditions during the Dekad. Pasture regeneration and animal's health conditions remains generally poor. (Figs. 1-4)
- The past rains have improved fodder for livestock, hence an increase in milk production and also availability of fresh vegetables.

CROP AND WEATHER REVIEW

(11th to 20th June, 2009)

Central Province and Nairobi Area:

Cool and cloudy mornings with light rains/ drizzle over high ground areas, was experienced over most places in Central province, Nairobi area and its environs during the Dekad. Generally most crops (cash and food crops) are doing well and corresponding to normal growth. The maize crop is at flowering stage, while the bean crop is at the maturity stage. (Fig1&2).

Eastern Province:

Cool and cloudy mornings with light rains/ drizzle over high ground areas, was experienced over most places bordering Mt. Kenya (Embu, Runyenjes, Chogoria, Chuka and Meru) during the Dekad. Most crops (cash and food crops) are doing well and corresponding to normal growth. The maize crop is at flowering stage, while the bean crop is at the maturity stage. (Fig1&2)

In the Southern Lowlands of the Eastern Province (Machakos, Makueni, Mwingi and Kitui districts), sunny conditions was experienced over most places during the Dekad. The maize and beans crops are experiencing water stress due to insufficient rainfall. (Fig1&2).

Coast Province:

Heavy rainfall was experienced over several places during the Dekad (Lamu, Malindi, Kwale and Voi). Maize crop is at flowering stages and is doing well with normal yields being expected. (Figs.1-3)

North Eastern Province:

Sunny and dry conditions dominated the region (Mandera, Moyale, Wajir and Garissa) during the Dekad. Pasture regeneration and the animal's health conditions remained generally poor, due insufficient rainfall and death of livestock is imminent due to inadequate pastures and water. (Figs.1, 2, 3 & 4)

Famine/hunger is still being experienced in most places and malnutrition levels are expected to be on rise.

Western Province:

Sunny conditions with light rains were experienced over over most places of this region (Kakamega, Bungoma, Busia) during the Dekad. Generally most crops (cash and food crops) are doing well and corresponding to normal growth. The maize crop is at flowering stage, while the bean crop is at the maturity stage and both are doing well and corresponding to normal growth. (Figs.1-3).

Nyanza and Central Rift Valley:

Light to moderate rainfall were experienced over several places in this regions (Kisumu, Nyamira, kisii Eldoret, Kitale, Kericho) during the Dekad. The maize and sorghum are at the flowering stages while the bean crop is at maturity. In areas around Kisumu crops have started wilting due to insufficient rainfall and poor yields being expected. (Figs.1-3).

In areas around Eldoret Kapsoya and Kitale the maize and bean crop are at the flowering stages and are wilting due to insufficient rainfall, with poor yields being expected. Adverse reports indicate the infestation of the maize crop by Maize Stock Borers and the bean crop by Aphids.

In areas around Kisii maize crop is at the flowering stage while the and bean crop is at harvesting stage and both are doing well with normal yields being expected. (Figs.1-3).

Northern and Southern Rift Valley:

Northerwestern Rift Valley (Lodwar, Lokitang) experienced generally sunny and dry conditions with light rains over few places during the Dekad. This has resulted in inadequate pastures and water sources for livestock, wildlife and human use leading to death of livestock and wildlife. (Figs.1-4).

Famine/hunger and its related impacts is still being experienced and malnutrition levels are still high.

The pastoral regions of Southern Rift Valley(Narok, Kajiado, Mara) experienced sunny and dry conditions resulting in inadequate water sources for livestock, wildlife and human use. Despite the rains in the previous Dekads, pastures regeneration has been gradual resulting in poor animals` health (both livestock and game). This situation has forced the pastolarists to drive their livestock in the National parks /game reserves posing unfair competention to the only available pastures for the wildlife.

For the farming community, wheat and maize crops are experiencing water stress due to inadequate rainfall.




Fig. 1: Actual Rainfall (11th to 20th June, 2009) in mm





Fig 2: Mean Temperatures (11th to 20th June, 2009) deg. Celsius



NDV_20090611

Clouds Bare Soil

Sparse Veg

Light Veg

📕 Medium Veg

Heavy Veg Water

Afr_tiny

FIG 3: DEKAD 17 Normalized Difference Vegetation Index (11th to 20th June, 2009)






FIG 4: DEKAD 16 Normalized Difference Vegetation Index (1st to 10th June, 2009))

EXPECTED WEATHER AND CROP CONDITIONS DURING THE NEXT 10 DAYS (21st - 30th June 2009).

During the next 10 days (21st – 30th June, 2009) the Western regions are expected to experience light to moderate rainfall over several places. Crops are expected to continue doing well and correspond to normal growth with normal yields being expected.

Nyanza and Central Rift Valley are expected to experience light to moderate rainfall over few places, poor crops performance are expected due to reduced rainfall with below normal yields being expected in areas around Kisumu, Kitale and Eldoret.

Central Highlands, Nairobi area and its environs, are expected to experience light rains over few places, with cool and cloudy mornings with light rains/ drizzle over high ground areas. Crops are expected to continue doing well and correspond to normal growth. The beans is expected to reach full maturity stages and with normal yields being expected.

In Eastern Province regions of Embu and Meru districts, bordering Mt Kenya, are expected to experience light rains over few places, with cool and cloudy mornings with light rains/ drizzle over high ground areas. Crops are expected to continue doing well and correspond to normal growth. The beans is expected to reach full maturity stages and with normal yields being expected.

In South-Eastern lowlands, generally sunny conditions are expected. Poor crop performances are expected due to insufficint rainfall.

The Coastal region is expected to experience moderate to heavy rainfall. Good crop performance is are expected due to increased rainfall.

In the pastoral regions and game parks/reserve of Northwestern, Northeastern, Northern Rift Valley, Southern Rift Valley and Southeastern Lowlands, sunny conditions are expected to prevail. Poor pastures regeneration and inadequate water sources for human and animal/wildlife use are expected. Poor animals` health both (livestock and game) are expected, due to inadequate pastures.

In summary, Crops are expected to continue doing well and correspond to normal growth in Western, parts of Nyanza, Central Rift Valley, Central Highlands and parts of Eastern province bordering Mt. Kenya. In the pastoral region and game parks/reserves poor pasture regeneration and water sources are expected.

This product should be used in conjunction with Kenya Meteorological Department weather forecasts. For more information, Contact

The Kenya Meteorological Department Agrometeorological Division Dagoretti Corner, Ngong Road, P.O. Box 30259, 00100 GPO, Nairobi Tel: +254 (0)20 3867880-7/3876957/3873682 Fax: +254 (0)20 3876955 E-mail: agromet@meteo.go.ke Website: www.meteo.go.ke

©2009 The Kenya Meteorological Department